

## Role of Anthropometric Dimensions of Human Body in Identifying Temperament in Traditional Persian Medicine

A. Vahedi<sup>1</sup>, M. Zamani<sup>1</sup>, M. Mojahedi(MD,PhD)<sup>\*2,3</sup>, S.A. Mozaffarpur(MD,PhD)<sup>2,3</sup>,  
R. Saghebi (MD,PhD)<sup>2,3</sup>, M.A. Mououdi (MSc)<sup>4</sup>

1. Student Research Committee, Babol University of Medical Sciences, Babol, I.R.Iran

2. Traditional Medicine and History of Medical Sciences Research Center, Babol University of Medical Sciences, Babol, I.R.Iran

3. Department of Traditional Medicine, Faculty of Traditional Medicine, Babol University of Medical Sciences, Babol, I.R.Iran

4. Department of Health, Faculty of Health, Mazandaran University of Medical Sciences, Sari, I.R.Iran

J Babol Univ Med Sci; 18(7); Jul 2016; PP: 24-33

Received: Feb 14<sup>th</sup> 2016, Revised: Mar 2<sup>th</sup> 2016, Accepted: Jun 1<sup>th</sup> 2016

### ABSTRACT

**BACKGROUND AND OBJECTIVE:** From the viewpoint of Traditional Persian Medicine (TPM), temperament of each person influences his physical and physiological properties such as body dimensions. The aim of this study is to review the reasons behind the diversity of human anthropometric measurements and their status in identifying temperament of people.

**METHODS:** In this descriptive study, we searched online databases such as Sid.ir, PubMed, Scopus, Magiran.com and Google Scholar for Persian key words such as "Anthropometry", "ergonomics" and "temperament" and their English equivalent. Authentic TPM books such as "The Canon of Medicine" by Avicenna, "Complete Book of the Medical Art" by al-Majusi, "al-Mansouri fi al-Tibb" (The book on medicine dedicated to al-Mansur) by Zakariya al-Razi, "Kholasa't ol Hikma" (summary of wisdom) by Aghili Khorasani, "Zakhireh kharazmshahi"(The treasure of Kharazm Shah) by Ismail Jurjani and "Bahr al-jawahir" (sea jewels) were also studied.

**FINDINGS:** Results of the study demonstrated that there is a direct relationship between weight gain, BMI and dimensions of soft tissue which are primarily signs of obesity and fat gain and cardiovascular diseases and diabetes. Since increase in the aforementioned indices can be a sign of coldness and wetness of temperament, one can argue that people with cold and wet temperament are more susceptible to such diseases. In references of TPM, temperament is mentioned as an agent that changes body dimensions and among the indices that identify temperament, "shape of organs" and "physique" is related to anatomic dimensions of body and obesity and thinness condition, receptively. Magnitude of chest and other organs is a sign of hotness; thinness is a sign of dryness; dominance of muscle tissue is a sign of hotness and wetness and dominance of adipose tissue is a sign of coldness and wetness of temperament.

**CONCLUSION:** According to the results of the present study, variety of anthropometric dimensions is related to genetic loci. Proving the hypothesis of relationship between anthropometric dimensions and temperament and relationship between temperament and genetic polymorphism in TPM requires more research. If the mentioned relationship is confirmed, the process of accessing standard tools for identifying temperament will be facilitated.

**KEY WORDS:** *Anthropometric, Ergonomics, Temperament, Traditional Medicine.*

### Please cite this article as follows:

Vahedi A, Zamani M, Mojahedi M, Mozaffarpur SA, Saghebi R, Mououdi MA. Role of Anthropometric Dimensions of Human Body in Identifying Temperament in Traditional Persian Medicine. J Babol Univ Med Sci. 2016;18(7):24-33.

\*Corresponding author: M. Mojahedi (PhD)

Address: Faculty of Traditional Medicine, Babol University of Medical Sciences, Ganjafrouz St, Babol, I.R.Iran

Tel: +98 11 32194730

E-mail: mortazamojahedy@gmail.com

## Introduction

Anthropometry, a subdivision of anthropology, measures dimensions and diameters of human body as a branch of anatomy science (1, 2). Although the main application of this science is in designing sites and equipments for humans, some standardized indices of this field such as thigh circumference, waist circumference and body mass index (BMI) are applicable in medicine. Most of the studies in this field are associated with identifying the relationship between anthropometric indices and occurrence or absence of some diseases (3-6).

Like many medical schools of ancient civilizations, Traditional Persian Medicine (TPM) has placed the basis of its health and illness diagnosis and health protection guidelines on individualized differences. This school of medicine has categorized the physical and mental differences in the human in the form of the keyword "Temperament" (called "Mizaj" in Persian language and TPM), investigated and presented its variations carefully (7, 8). From the viewpoint of TPM, temperament is the result of interaction between body organs; it is not identical in any two people in the world, so any member of human society has a unique temperament (9). TPM scholars have used different indices to identify temperament. Avicenna, the prominent Iranian physician, has categorized the indices that identify temperament into ten groups and explained their role in identifying temperament in full detail (10).

Among the ten aforementioned indices, two indices of "shape of organs" and "physique" (general appearance of the body) pertain to body dimensions. The purpose of this study is to investigate recent researches about the reason of diversity in anthropometric dimensions of different races and particularly members of one race as well as summarizing the opinions of TPM scholars about application of anthropometric dimensions in identifying temperament. If there is a probable relationship between varieties of anthropometric dimensions and differences in temperament, the process of explaining physiologic basics of temperament in TPM will be facilitated.

## Methods

In this descriptive study, we searched online databases such as Sid.ir, PubMed, Scopus, Magiran.com and Google Scholar for Persian key

words such as "Anthropometry", "ergonomics" and "temperament" and their English equivalent. Authentic TPM references such as "The Canon of Medicine" by Avicenna, "Complete Book of the Medical Art" by al-Majusi, "al-Mansouri fi al-Tibb" (The book on medicine dedicated to al-Mansur) by Zakariya al-Razi, "Kholasa't ol Hikma" (summary of wisdom) by Aghili Khorasani, "Zakhireh kharazmshahi" (The treasure of Kharazm Shah) by Ismail Jurjani and "Bahr al-jawahir" (sea jewels) were also studied.

## Results

After investigating authentic databases, 54 articles and 8 books were ultimately used in the study.

**A. Variety of anthropometric indices among different races:** Science of ergonomics has elicited and standardized various indices to measure anthropometric dimensions. For instance, indices of height, weight, sole length, shoulder width, chest width, waist-to-hip ratio (WHR) and thigh circumference are the most commonly used anthropometric indices. Studies have demonstrated that anthropometric indices are diverse among different ethnic groups and races.

In other words, no two people in this world are alike, not even identical twins (11). Results of various studies in different countries show that length of organs is diverse among people (12, 13). In the study of Sanli et al. on 155 adult Turkish men and women, mean sole length, palm length and height of men was reported more than women (14). In Iran, study of Meshkdanian et al. on 300 Qazvin (an Iranian city) natives demonstrated that mean upper limb length, arm length, forearm length, hand length and hand width of men is more than women (15). In the aforementioned study, a comparison with the study of Holliday et al. revealed that mean forearm length of Qazvini men was more than Pakistani and Hindi men and the same index in Qazvini women was more than Pakistani and German women (15, 16).

Joneidi et al. have investigated 36 anthropometric indices among 6 Iranian ethnic groups in their study and reported significant differences in some indices; anthropometric indices of each ethnic group have been reported as a range of numbers (table 1) (17). Results of different studies suggest that ethnicity and inheritance can influence face shapes of people (18-21). Findings of Jahanshahi et al. in the north of Iran have demonstrated that women's face is generally

Euryprosopic and men's face is generally Mesoprosopic among Fars and Turkmen races (19). In the study of Ghosh et al. on Santhal tribe in India, it has been found that women's face is generally Hypereuryprosopic and men's face is generally Euryprosopic (22). Body mass index (BMI) is different among people in European countries, American ethnic groups and various races in Asian and African countries (23-27). Although all measurements of anthropometric indices are reported as a range of numbers, no detailed study has been found to report difference in anthropometric indices of people in a particular race to the extent of results of this research.

**Table 1. Anthropometric dimensions of some Iranian Turk male workers (cm) (17)**

| Variable           | Mean±SD | 5 <sup>th</sup> percentile | 95 <sup>th</sup> percentile |
|--------------------|---------|----------------------------|-----------------------------|
| Height             | 169±10  | 153                        | 185                         |
| Length of knee/hip | 57±3    | 53                         | 62                          |
| Shoulder width     | 44±4    | 38                         | 50                          |
| Chest depth        | 23±3    | 20                         | 28                          |
| Upper limb length  | 78±7    | 68                         | 88                          |
| Hand width         | 8±1     | 7                          | 9                           |
| Shoulder length    | 34±3    | 28                         | 39                          |
| Hip width          | 40±3    | 35                         | 45                          |

**B. Factors that cause variety in anthropometric dimensions:** Study of Pietiläinen et al. and Silventoinen et al. have demonstrated that genetics and environmental factors can influence changes in body dimensions from childhood to early adulthood (28, 29). Study of Slemenda et al. has demonstrated that various patterns of ossification can be inherited and genetic factors can influence familial resemblance of bone mass in the entire skeletal system and in all ages (30). On the other hand, it was observed that inheritance and genetics plays a role in obesity of children and adults. For instance, study of Ochoa et al. demonstrated that synergistic interaction between PPAR $\gamma$ 2 and ADR $\beta$ 3 genes increases obesity risk up to 20 times in children and adolescents that carry variants of these two genes (31).

Scuteri et al. have shown that variants of FTO and PFKP genes are associated with increase in BMI, hip circumference and weight of carriers (32). Herbet et al. have also indicated that genetic variants near INSIG2 gene are associated with obesity (33). Soranzo et al. have indicated in their study that 17 gene loci are associated with height and CATSPER4 and TMED10 are the major relevant loci among them. GPR126,

LCORL and PRKG2 are associated with body length, hip axis length (HAL) and thigh length, respectively. Nucleotide differences in these loci cause differences in skeletal dimensions among people (34). In a study of 146 20-83 years old men, Need et al. have found that presence of polymorphism in vitamin D receptor (VDR) gene is the factor that causes diversity in bone density and bone dimensions among men. People with BB genotype are likely to have bigger bone dimensions than other people (35).

Results of various studies demonstrate that diversity in nutrition and activities of individuals can influence their anthropometric indices such as height, weight and BMI (36-38). Studies of Elshibly et al. have demonstrated that there is a significant relationship between maternal lean body mass and weight, height and circumference of baby. Moreover, it has been observed that delivery order and mother's height are associated with risk of low birthweight and these two parameters have inverse relationship with the risk of underweight (39, 40).

**C. Application of anthropometric indices:** Anthropometry, as a major branch of ergonomics, plays a key role in advancing the goals of ergonomics by studying body dimensions and helping them to adapt to work tools and equipments (41, 42). Although the main application of anthropometry is in designing sites and equipments, some standardized indices of this field such as thigh circumference, waist circumference and body mass index (BMI) are applicable in medicine to detect physiological differences and their relationship with illnesses. Previous studies have demonstrated that there is a relationship between general power of organs and body and some of the anthropometric indices (43-45).

Study of Martin et al. have shown that there is a direct relationship between hand grip strength and volume of forearm muscles and upper arm muscle (46). Various studies have indicated that anthropometric indices such as height and weight can affect menstruation (47-49). In a study of Qaravi et al. on 100 female university students in Gorgan, Iran, it was found that there is a significant direct relationship between girls' height and age of first menstruation (50). Regarding cardiovascular diseases, findings of Heitmann et al. have demonstrated that smaller thighs mean greater risk of cardiovascular diseases (3). Moreover, findings of Debette et al. and Kim et al. have demonstrated that reduction in the size of calf is associated with greater risk of cardiovascular diseases

(4, 5). Findings of Li et al. and Chuang et al. have demonstrated that index of wrist circumference is more associated with type 2 diabetes than other indices (51, 52). Nowadays in many developed countries, anthropometric indices are used in different surgeries such as plastic surgery, oral and maxillofacial surgery and brain surgery (53-55).

In a study of 400 native boys and girls of Esfahan, Alavi et al. have shown that anthropometric dimensions of skull and face among people of Esfahan is different from the same indices among people of Canada and difference in anthropometric dimensions in various races requires normal anthropometric index that is appropriate for that race in surgeries and tools for everyday life (56).

**D. Reasons of variety in anthropometric indices and their application in traditional medicine:** TPM has categorized the physical and mental differences of members of human societies in the form of the keyword "Temperament" and has studied its changes. From the viewpoint of this school, temperament is the result of interaction between body organs; it is not identical in any two people in the world and any member of human society has a unique temperament (57-59). This temperament diversity is evaluated in the form of two different ranges of quality: hotness and coldness, and dryness and wetness (10,60). In fact, temperament is a consequent of four qualities (hotness, coldness, dryness and wetness) of different components of a compound like human body (9,10, 60-62).

**1. Status of temperament in TPM:** Most healthcare directives of this school, which are presented under the title "Health Protection Measures", are set based on people's temperament and each healthcare directive is different from one person to another based on the type of temperament. On the other hand, illness diagnosis and treatment are done according to the type of temperament. One the important principles of prescribing medicine in this school is careful consideration of temperament of patient, temperament of patient's organs, temperament of the type of illness and finally temperament of the drug (58, 63).

Resources of this study have indicated that each person comes to this world based on a specific model of temperament. Avicenna has proposed temperament modification in principle of vigor and stated that the primary temperament of each person can change as people get older and their eating habits and other occupational and personal habits change (10).

Resources of this study have categorized temperament into nine groups with details: hot, cold, wet, dry, hot and wet, hot and dry, cold and wet, cold and dry and moderate (10).

On the other hand, temperament influences physical, mental and physiological characteristics. Hence, in order to identify a person's temperament and their status in one of the nine groups it is necessary to examine all the related indices to identify their temperament (9, 10, 60-62).

**2. Indices that identify temperament:** Since many mental and physical indices need to be considered to identify temperament, Avicenna categorized these indices into ten groups under the title "ten materials" as follows: 1. Body sense. 2. Obesity and thinness (physique) 3. Hair. 4. Skin color. 5. Body dimensions (shape of organs). 6. Impressionability speed. 7. Sleep-wake states. 8. Actions. 9. Condition of fecal material. 10. Mental interactions (10). Of the aforementioned indices, shape of organs and physique were the focus of this study.

**2.1 Shape of organs:** Shape refers to form and figure of a thing (64). From the viewpoint of Avicenna, size of organs, joints and chest represents the general shape of organs. Magnitude of organs is a sign of hotness and smallness of organs indicates coldness of temperament. Hot temperament causes magnitude of chest and organs, width and salience of superficial veins and magnitude of muscles and their proximity to joints; such indices are rarely observed in people with cold temperament (10). Jurjani sees wide chest and large veins as signs of hotness and lack of these characteristics as coldness and the median condition as a sign of moderation (9).

Ahwazi considers strength of organs as a sign of hot temperament and magnitude of chest (particularly along with smallness of head) a sign of hot heart and considers the opposite symptoms as a sign of coldness of heart temperament (60). Razi notes this index as one the main indices that identify temperament and mentions magnitude of nostrils and ducts in people with hot temperament and mentions narrowness and thinness of them in people with cold temperament. He mentions lack of salience of joints as a sign of wet temperament, salience of joints as a sign of dry temperament and salience of tendons, bones and joints as a sign of hot and dry temperament. He sees salient larynx, long neck and long nose as signs of cold temperament and magnitude and salience of eyes, width of nose, beef cheek, little facial hair and thin

nails as signs of wet temperament. He also sees shortness and thickness of fingers as a sign of coldness and wetness of temperament, respectively (61).

**2.2 Physique:** In TPM, physique refers to obesity and thinness status and weakness and stiffness of adipose tissue and muscles and divides people into three groups: fat people, thin people and moderate people (62). Increase in adipose tissue and muscles compared with the average level of community is a sign of wetness, decrease in adipose tissue is a sign of dryness and a moderate status between obesity and thinness is

a sign of moderate temperament. A fat body with dominance of adipose is a sign of coldness and wetness, a muscular body is a sign of hotness and wetness and thinness is usually a sign of dryness and is not an indicator of hotness or coldness alone. Nevertheless, cold and dry people are usually thinner than hot and dry people; although hot and dry people are thin, they have an acceptable muscular tissue, while cold and dry people lack a notable muscular tissue (table 2) (58, 60).

**Table 2. Shape of organs and physique as two indices to define temperament in Iranian traditional resources**

| Index           | Wetness                                 | Dryness                         | Hotness                          | Coldness                              |
|-----------------|-----------------------------------------|---------------------------------|----------------------------------|---------------------------------------|
| Physique        | Dominance of adipose or muscle          | Thinness                        | Dominance of muscle over adipose | Dominance of adipose/Extreme thinness |
| Shape of organs | Softness of organs (in movements)       | Long neck                       | Magnitude of chest               | Smallness of chest                    |
|                 | Lack of salience of joints              | Salience of joints              | Magnitude of hands and legs      | Smallness of hands and legs           |
|                 | Lack of salience of laryngeal cartilage | Salience of laryngeal cartilage | Magnitude of muscles             | Smallness of muscles                  |
|                 | Wide nose                               | Thin and long nose              |                                  | Shortness of fingers                  |
|                 | Magnitude and salience of eyes          |                                 |                                  |                                       |
|                 | Beef cheek                              |                                 |                                  |                                       |
|                 | Thickness of fingers                    |                                 |                                  |                                       |

## Discussion

According to the results of this study, diversity of anthropometric dimensions exists among various races and also the people of one race. Although no study has been reported to investigate the differences among people of one race, diversity of anthropometric dimensions among homogeneous people has led to reportage of a range of anthropometric indices in homogeneous people of one race. Since TPM, based on the basic principle of temperament, has special attention to physical and biological differences of humans and uses them to diagnose and treat diseases and suggest health care directives, having access to practical methods of identifying temperament can help health promotion in society.

According to the fact that two practical indices of identifying temperament-shape of organs and physique-refer to anthropometric dimensions and characteristics of human body based on theoretical principles of traditional medicine, if the relationship between these dimensions and temperament of people is proved in future studies, it is possible to convert one of the qualitative indices of identifying temperament to a quantitative index by measuring the aforementioned

dimensions. Based on the results of this study, there is a direct relationship between weight gain, BMI and dimensions of soft tissue which are primarily signs of obesity and fat gain and cardiovascular diseases and diabetes. Since increase in the aforementioned indices can be a sign of coldness and wetness of temperament, one can argue that people with cold and wet temperament are more susceptible to such diseases. Moreover, since the risk of such diseases among thin people is lower than fat people and thinness is a sign of wet temperament, one can propose a hypothesis to assert that people with wet temperament are less susceptible to these diseases.

It is suggested that the relationship between people's temperament and susceptibility to disease be investigated in future studies. Furthermore, according to the results of this study, majority of the studies regarding relationship between diseases and anthropometric dimensions have investigated the relationship between soft tissue and aforementioned diseases and not many detailed and adequate studies have been conducted to investigate the relationship between osseous dimensions of body and various

diseases. Hence, it is suggested that more attention be paid to such issues in future studies. Since body dimension is one the indices that identifies temperament in TPM, any factor that influences these dimensions is theoretically a factor that can change people's temperament.

According to the fact that genetic differences of people can play a key role in diversity of their anthropometric indices, one can hypothesize that temperament in TPM might be closely related to some genetic properties. Based on this hypothesis one can argue that genes such as FTO and PFKP which cause obesity, act more effectively among people with cold and wet temperament. In other words, there may be a significant relationship between genetic polymorphism and temperament variety.

Based on the same hypothesis, one can expect to find a significant relationship between polymorphism of ossification loci (such as PRKG2, GPR126 and LCORL), which increases osseous dimensions, and hot temperament in TPM. To prove these hypotheses, having access to standard tools for identifying temperament of people and studying genetic

differences of these people is necessary. It is suggested that the relationship between different anthropometric indices of traditional medicine and various temperaments be investigated in future studies. If the aforementioned relationship is proven and proper cutoff points are extracted for related anthropometric indices, it will be a huge step forward in quantifying qualitative indices of identifying temperament and the process of accessing genetic basics of temperament will be accelerated. If these cases are studied in ergonomics science, a new approach is formed that offers anthropometric dimensions of each society based on temperamental differences of people and identifies workplace requirements, personal tools and equipments, and environmental conditions of workplace and living place.

### Acknowledgments

Hereby, we would like to thank the authorities and colleagues in Faculty of Traditional Medicine of Babol University of Medical Sciences for their unsparing cooperation.

## References

- 1.Lasker G. The place of anthropometry in human biology; Anthropometry: the individual and the population. London: Cambridge University Press; 1994.p.14:1.
- 2.Mououdi MA. Anthropometric engineering, 1<sup>st</sup> Tehran: Tehran university pub;1996.p. 20-35. [In Persian]
- 3.Heitmann BL, Frederiksen P. Thigh circumference and risk of heart disease and premature death: prospective cohort study. *Bmj*; 2009.p.339.
- 4.Debette S, Leone N, Courbon D, Gariépy J, Tzourio C, Dartigues J-F, et al. Calf circumference is inversely associated with carotid plaques. *Stroke*. 2008;39(11):2958-65.
- 5.Kim S-K, Choi YJ, Huh BW, Kim C-S, Park SW, Lee EJ, et al. Ratio of waist-to-calf circumference and carotid atherosclerosis in korean patients with type 2 diabetes. *Diabetes Care*. 2011;34(9):2067-71.
- 6.Peymani P, Heydari ST, Ahmadi SM, Sarikhani Y, Joulaei H, Moghadami M, et al. The prevalence of high blood pressure and its relationship with anthropometric indicators; a population based study in fars province, IR Iran. *Int Cardiovasc Res J*. 2012;6(2):40-5.
- 7.Low K, Ang S-L. The foundation of traditional chinese medicine. *J Chinese Med, Scie Res*. 2010;1(3):84-90.
- 8.Naseri M. The school of traditional Iranian medicine: The definition, origin and advantages. *Iran J Pharm Res*. 2010;3(Suppl 2):20.
- 9.Jorjani SE. Zakhireh Kharazmshahi. Moharrari Mr, editor. Tehran: The academi of Medical Science Islamic Republic of Iran 2002. [In Persian]
- 10.Ibn Sina (Avicenna) H. Canon of Medicine. Beirut-Lebanon. Alaalami Library;2005.p: 31-9, 170-7. [In Arabic]
- 11.Shokoohi H, Khoshroo M. Fitness measurement of the combat troops to prevent the skeleton and muscular disorders due to inappropriate design of the military equipments. *Ann Mil Health Sci Res*. 2011;9(3):172-7. [In Persian]
- 12.Ahmed AA. Estimation of stature from the upper limb measurements of Sudanese adults. *Forensic Sci Int*. 2013;228(1-3):178.
- 13.Krishan K, Sharma A. Estimation of stature from dimensions of hands and feet in a North Indian population. *J Forensic Leg Med*. 2007;14(6):327-32.
- 14.Sanli SG, Kizilkanat ED, Boyan N, Ozsahin ET, Bozkir MG, Soames R, et al. Stature estimation based on hand length and foot length. *Clin Anat*. 2005;18(8):589-96.
- 15.Moshkdanian Gh, Moghani Ghoroghi F, Shiasi M, Hassanzadeh G, Alaghebandha N, Dehbashipour A, et al. Anthropometric characteristics of upper limb in Iranian and Pakistani subjects. *J Gorgan Uni Med Sci*. 2014;16(3):80-5.[In Persian].
- 16.Holliday TW, Ruff CB. Relative variation in human proximal and distal limb segment lengths. *Am J Phys Anthropol*. 2001;116(1):26-33.
- 17.Joneidi A, Sadeghi F. A survey on static anthropometric dimensions in 20-60year workers of six Iranian nations. *J Health Administrat*. 2008;11(32):11-24. [In Persian]
- 18.Golalipour M, Jahanshahi M, Haidari K. The variation of head and face shapes in female newborns in the South-East of the Caspian Sea (Iran-Gorgan). *Eur J Anat*. 2005;9(2):95-8.
- 19.Jahanshahi M, Golalipour M, Heidari K. The effect of ethnicity on facial anthropometry in Northern Iran. *Singapore Med J*. 2008;49(11):940-3.
- 20.Farkas LG, Katic MJ, Forrest CR, Alt KW,Bagic I,Baltadjiev G,et al. International anthropometric study of facial morphology in various ethnic groups/races. *J Craniofac Surg*. 2005;16(4):615-46.
- 21.Farahvash MR, Yegane R, Khorasani G, Ahmadi M, Farahvash B. Anthropometric analysis of faces and heads of 18 to 30-year old Persians and its comparison with Caucasian ethnic group. *Tehran Univ Med J*. 2011;69(6):359-65. [In Persian]

- 22.Ghosh S, Malik SL. Sex differences in body size and shape among Santhals of West Bengal. *Anthropol.* 2007;9(2):143-9.
- 23.Goh VH, Tain C, Tong TY, Mok HP, Wong M. Are BMI and other anthropometric measures appropriate as indices for obesity? A study in an Asian population. *J Lipid Res.* 2004;45(10):1892-8.
- 24.Wu H,Zhu K,Shah M,Lipnick RJ,Kao TC. BMI and Cervical Cancer Screening among White, African-American, and Hispanic Women in the United States. *Obesity.* 2006;14(3):526-7.
- 25.Ramel A, Halldorsson TI, Tryggvadottir EA, Martinez JA, Kiely M, Bandarra NM, et al. Relationship between BMI and body fatness in three European countries. *Eur J Clin Nutr.* 2013;67(3):254-8.
- 26.Wee CC, Phillips RS, McCarthy EP. BMI and Cervical Cancer Screening among White, African-American, and Hispanic Women in the United States. *Obes Res.* 2005;13(7):1275-80.
- 27.Yates A, Edman J, Aruguete M. Ethnic differences in BMI and body/self-dissatisfaction among Whites, Asian subgroups, Pacific Islanders, and African-Americans. *J Adolesc Health.* 2004;34(4):300-7.
- 28.Pietiläinen KH, Kaprio J, Räsänen M, Rissanen A, Rose RJ. Genetic and environmental influences on the tracking of body size from birth to early adulthood. *Obes Res.* 2002;10(9):875-84.
- 29.Silventoinen K, Pietiläinen K, Tynelius P, Sørensen T, Kaprio J, Rasmussen F. Genetic and environmental factors in relative weight from birth to age 18: the Swedish young male twins study. *Int J Obes (Lond).* 2007;31(4):615-21.
- 30.Slemenda CW, Christian JC, Williams CJ, Norton JA, Johnston CC. Genetic determinants of bone mass in adult women: a reevaluation of the twin model and the potential importance of gene interaction on heritability estimates. *J Bone Miner Res.* 1991;6(6):561-7.
- 31.Ochoa M, Marti A, Azcona C, Chueca M, Oyarzabal M, Pelach R, et al. Gene-gene interaction between PPAR $\gamma$ 2 and ADR $\beta$ 3 increases obesity risk in children and adolescents. *Int J Obes (Lond).* 2004;28:37-41.
- 32.Scuteri A, Sanna S, Chen W-M, Uda M, Albai G, Strait J, et al. Genome-wide association scan shows genetic variants in the FTO gene are associated with obesity-related traits. *Plos Genet.* 2007;3(7):115.
- 33.Herbert A, Gerry NP, McQueen MB, Heid IM, Pfeufer A, Illig T, et al. A common genetic variant is associated with adult and childhood obesity. *Science.* 2006;312(5771):279-83.
- 34.Soranzo N, Rivadeneira F, Chinappen-Horsley U, Malkina I, Richards JB, Hammond N, et al. Meta-analysis of genome-wide scans for human adult stature identifies novel Loci and associations with measures of skeletal frame size. *PLoS Genet.* 2009;5(4):e1000445-e.
- 35.Need A, Horowitz M, Stiliano A, Scopacasa F, Morris H, Chatterton B. Vitamin D receptor genotypes are related to bone size and bone density in men. *Eur J Clin Invest.* 1996;26(9):793-6.
- 36.Truel M, Rivera J, Habichat JP, Martorell R. Differential response to early nutrition supplementation: long-term effects on height at adolescence. *Inter J Epidemiol.* 1995;24(2):404-12.
- 37.Mazloom Z, Kazemy F, Tabatabai S, Ansar H. Comparison of the effect of low-glycemic index versus low-fat diet on body fat and waist-hip ratio in obese women. *J Gorgan Univ Med Sci.* 2009;11(1):33-8.
- 38.Kuhle CL, Steffen MW, Anderson PJ, Murad MH. Effect of exercise on anthropometric measures and serum lipids in older individuals: a systematic review and meta-analysis. *BMJ Open.* 2014;4(6):005283.
- 39.Elshibly EM, Schmalisch G. The effect of maternal anthropometric characteristics and social factors on gestational age and birth weight in Sudanese newborn infants. *BMC Public Health.* 2008;8(1):244.
- 40.Elshibly EM, Schmalisch G. Relationship between maternal and newborn anthropometric measurements in Sudan. *Pediatr Int.* 2009;51(3):326-31.
- 41.Pheasant S, Haslegrave CM. *Bodyspace: Anthropometry, ergonomics and the design of work.* 3ed:CRC Press; 2005.

42. Varmazyar S, Ghalehnovi M, Amani Z, F M, Aivazloo T, Inanloo F, et al. School Desk and Chair Design Based on High School Female Students Anthropometry Qazvin, Iran 2007 to 2008. *Qom Univ Med Sci J.* 2008;2(3):39-45. [In Persian]
43. Aghazadeh F, Lee K, Waikar A. Impact of anthropometric and personal variables on grip strength. *J Hum Ergol (Tokyo).* 1993;22(2):75-81.
44. Clerke AM, Clerke JP, Adams RD. Effects of hand shape on maximal isometric grip strength and its reliability in teenagers. *J Hand Ther.* 2005;18(1):19-29.
45. Link L, Lukens S, Bush MA. Spherical grip strength in children 3 to 6 years of age. *Am J Occup Ther.* 1995;49(4):318-26.
46. Martin S, Neale G, Elia M. Factors affecting maximal momentary grip strength. *Hum nutr Clin Nutr.* 1985;39(2):137-47.
47. Ayatollahi S, Dowlatbadi E, Ayatollahi S. Age at menarche in Iran. *Ann Hum Biol.* 2002;29(4):355-62.
48. Lee SE, Yang JY, Lee JH, Kim HW, Kim HS, Lee HJ, et al. Relationship of age at menarche on anthropometric index and menstrual irregularity in late adolescent girls in Seoul. *Ann Pediatr Endocrinol Metab.* 2013;18(3):116-21.
49. Zhu H, Pan H, Zhang D, Wu Q, Zhang K, Li M, et al. Effect of bodyweight on the onset of puberty of female children and adolescents. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao.* 2010;32(1):25-8.
50. Gharravi AM, Gharravi S, Marjani A, Moradi A, Golalipour MJ. Correlation of age at menarche and height in Iranian student girls living in Gorgan-Northeast of Iran. *J Pak Med Assoc.* 2008;58(8):426-9.
51. Li C, Ford ES, Zhao G, Kahn HS, Mokdad AH. Waist-to-thigh ratio and diabetes among US adults: the Third National Health and Nutrition Examination Survey. *Diabetes Res Clin Pract.* 2010;89(1):79-87.
52. Chuang Y-C, Hsu K-H, Hwang C-J, Hu P-M, Lin T-M, Chiou W-K. Waist-to-thigh ratio can also be a better indicator associated with type 2 diabetes than traditional anthropometrical measurements in Taiwan population. *Ann Epidemiol.* 2006;16(5):321-31.
53. Christofides EA, Steinmann ME. A novel anthropometric chart for craniofacial surgery. *J Craniofac Surg.* 2010;21(2):352-7.
54. Doddi NM, Eccles R. The role of anthropometric measurements in nasal surgery and research: a systematic review. *Clin Otolaryngol.* 2010;35(4):277-83.
55. Farkas L, Kolar J. Anthropometric guidelines in cranio-orbital surgery. *Clin Plast Surg.* 1987;14(1):1-16.
56. Alavi S, Safari A. An investigation on facial and cranial anthropometric parameters among Isfahan Young adults. *J Dent Med.* 2003;16(1):19-28. [In Persian]
57. Yousefifard M, Parviz M, Hosseini M, Ebadiani M, Keshavarz M. Mizaj; past, present and future. *Physiol Pharmacol.* 2013;16(4):328-39. [In Persian]
58. Mojahedi M, Naseri M, Majdzadeh R, Keshavarz M, Ebadiani M, Nazem E, et al. A Review on Identification Mizaj (Temperament) Indices in Iranian Traditional Medicine (ITM). *Quart J Med His.* 2012;4(12):37-76. [In Persian]
59. Behmanesh E, Mojahedi M, Saghebi R, Ayati MH, Bahrami Taghanaki HR, Allameh H, et al. Comparison between the "Theory of Arkan"(Four Elements) in Traditional Persian Medicine and the "Theory of Five Elements" in Traditional Chinese Medicine. *Res Hist Med.* 2015;4(2):73-86. [In Persian]
60. Ahvazi AEA. *Kamil al-Sinaā al Tibbiya.* [Qom-Iran: Jallaleddin 2008. [In Arabic]
61. Rhazes MZ. *Al-Mansuri Fi At-tebb.* In: Al-Siddiky HA-B, editor. Kuwait: Institue of Arab manuscripts 1987; pp: 79-92. [In Arabic]
62. Aghili Khorasani shirazi MH. *Kholassat Al-Hekmah (The Principal's of Traditional Iranian Medicine),* Nazem E. Qom: Esmaeilian;2006.p: 482-7,895-902. [In Persian]

- 63.Mojahedi M, Naseri M, Majdzadeh R, Keshavarz M, Ebadini M, Nazem E, et al. Reliability and validity assessment of Mizaj questionnaire: a novel self-report scale in Iranian traditional medicine. Iran Red Crescent Med J. 2014;16(3):e15924.
- 64.Heravi M. Bahr-al-javaher, Qom: Jallaleddin; 2008. [In Persian]