

The Relationship between Atopic Dermatitis and Premenstrual Syndrome in Women Referred to the Skin Clinic of Hamadan

S.Z. Masoumi (PhD)¹, F. Kazemi (PhD)², M.R. Sobhan (MD)³, S.R. Oliaei (DDS)⁴,
N. Hoseini (BSc)^{5*}, Gh. Roshanaie (PhD)⁶

1. Research Center of Mother & Child Care, Faculty of Nursing and Midwifery, Hamadan University of Medical Sciences, Hamadan, I.R.Iran.

2. Department of Midwifery, Faculty of Nursing and Midwifery, Hamadan University of Medical Sciences, Hamadan, I.R.Iran.

3. Department of Dermatology, Faculty of Medicine, Hamadan University of Medical Sciences, Hamadan, I.R.Iran.

4. Faculty of Dentistry, Hamadan University of Medical Sciences, Hamadan, I.R.Iran.

5. Department of Public Health, Faculty of Public Health, Hamadan University of Medical Sciences, Hamadan, I.R.Iran.

6. Research Center of Non-Communicable Disease Modeling, Hamadan University of Medical Sciences, Hamadan, I.R.Iran

J Babol Univ Med Sci; 20(5); May 2018; PP: 54-60

Received: Oct 21st 2017, Revised: Feb 17th 2018, Accepted: Mar 26th 2018.

ABSTRACT

BACKGROUND AND OBJECTIVE: One of the factors affecting dermatophyte is hormonal factors, in which atopic dermatitis is more common in adult women. Since one of the psychotic problems regarding reproductive performance of women is premenstrual syndrome, in this study, the association between premenstrual syndrome and atopic dermatitis in reproductive age women was investigated.

METHODS: This case-control study was performed on 82 women aged between 15 and 49 years old in two groups of atopic dermatitis referring to Skin Clinic of Sina Hospital in Tehran and control group including healthy women for atopic dermatitis. Demographic questionnaire and (PSST) Premenstrual Symptoms Screening Test (with score 42-0) were completed and evaluated by both groups.

FINDINGS: The mean age of subjects in the case group was 17.17 ± 4.45 and in the control group was 27.29 ± 8.03 years. The results showed by controlling the possible confounding factors as age, occupation and marital status the probability of having moderate or severe premenstrual syndrome in patients with atopic dermatitis is 13.26 (OR= 13.26, CI-95%: 3.31, 53.07), as compared with healthy subjects.

CONCLUSION: The results showed that there is a strong relationship between moderate to severe menstrual syndrome and atopic dermatitis in women.

KEY WORDS: *Atopic Dermatitis, Premenstrual Syndrome, Case-Control Study, Women.*

Please cite this article as follows:

Masoumi SZ, Kazemi F, Sobhan MR, Oliaei SR, Hoseini N, Roshanaie Gh. The Relationship between Atopic Dermatitis and Premenstrual Syndrome in Women Referred to the Skin Clinic of Hamadan. J Babol Univ Med Sci. 2018;20(5):54-60.

* Corresponding Author: N. Hoseini (BSc)

Address: Pazhouhesh crossroad, Shahid Fahmideh Blvd, Facing the Mardom Park, Hamadan University of Medical Sciences, Hamadan, I.R.Iran.

Tel: +98 81 38380150

Email: narmin.hosseini@gmail.com

Introduction

Premenstrual syndrome (PMS) is one of the psychotic problems regarding to reproductive function of women associated with behavioral and emotional disturbances such as anger, anxiety and irritability (1). The most common age of women with the syndrome is 20-35 years of age (3, 2).

Premenstrual syndrome in the 10th International Classification of Mental, Behavioral and Developmental Disorders is categorized as a component of women's medical disorders and not psychiatric disorders, which indicate the temporary state of impairment at certain times (premenstrual) (5, 4). Nowadays psychological disorders have a special place in psychiatry, and each passing day, the relationship between physical illness and human psyche is becoming more and more evident, which also requires more emphasis on mental health for the better treatment of these diseases.

Among the various medical disciplines, the skin line has the highest number of psychotic patients who are referred to dermatologists due to a lack of familiarity with psychiatric services or a reluctance to refer to a psychiatrist or psychologist. Of these patients, only a small percentage is presented by the dermatologist for psychological treatment, the rest of them are only examined and treated in terms of skin complaints, and they usually do not endorse commonly used treatments (4, 2), which causes their problems not to be fully addressed and refer to many physicians and treatment centers for complete treatment, which imposes a heavy burden on the individual and the community (6). The findings have shown that hormonal factors are effective on dermatophytes and atopic dermatitis is more common in adult women (8, 7).

Atopic dermatitis is a chronic inflammatory disease with an outbreak of about 17-8% in adults under the age of 60 years (9) that is chronic in severity and healing periods and is diagnosed with cutaneous eczema lesions (10) and can have a genetic background (12, 11). In patients with eczema, lesions (papules and plaques) are inflamed as periodic.

The lesions may be intensified by skin irritation, infection, stress, and other factors (13). The difference of constitutional atopic dermatitis form of other dermatitis is in the level of immunoglobulin E, which is normal in atopic dermatitis, as well as these individuals have no specific IgE and no immediate reaction symptoms (14-16). According to previous studies, about 25% of women experienced atopic dermatitis

improvement during pregnancy and more than 50% of women with atopic dermatitis had a deterioration during this period (18,17). In addition, women with atopic dermatitis often show worsening of skin symptoms during the menstrual period (19).

Timonen and colleagues showed that the relationship between atopic and depressive disorders is limited to women, and this relationship has not been found among men. They also stated that female hormones may be a contributing factor to the relationship between depression and atopic disorders, such as eczema (21, 20). The basis of these reports is the relationship between the number of pregnancies and the gestational age and atopic disorders, and the relationship between multiple atopies and depression (23, 22).

Relationships between premenstrual syndrome and various diseases including allergies, asthma, eczema, allergic rhinitis, acne, epilepsy and migraine have been studied (24). Considering the importance of psychosomatic diseases and premenstrual syndrome and its psychological symptoms such as depression, anxiety and emotional tension that are common in 90% of women, although the association between these symptoms and the premenstrual syndrome itself with atopic dermatitis was studied, but a similar study has not been conducted in these studies in Iran; since due to the relationship between hormonal changes and its effect on the exacerbation of skin diseases, a large amount of treatment costs are spent on temporary treatment of the disease.

Therefore, in this study, the association between premenstrual syndrome and atopic dermatitis in women of reproductive age were studied to minimized treatment costs of the disease as much as possible.

Methods

This case-control study was done on the women of reproductive age referred to the skin care clinic of Sina Hospital, Hamedan, in 2016 and in a 6-month period after approval by the Ethics Committee of Hamedan University of Medical Sciences with registration code of IR.UMSHA.REC.1394.558.

Considering the statistical power of 0.95, the error level of 0.05, the sample size ratio in the case group to the control (r) is equal to one, and the maximum difference between the two proportions that these two maximal sample sizes are presented according to previous studies. In this regard, for each case and

control group, there were 41 individuals and 82 individuals in total. Women aged 15- 49 years without mental illness and ability to understand and respond to the questionnaire, lack of smoking and alcohol was included in the study.

In the case of pregnancy, irregular menstrual cycle, the use of hormonal contraceptives and neurological drug consumption, and infectious dermatitis, such as fungal diseases were excluded. Among women referring to the skin clinic, if they were willing to participate in the study, written consent was obtained and then referred to the dermatologist for confirmation or rejection of atopic dermatitis. The method of diagnosis of atopic dermatitis was based on Hanifin and Rajka criteria. According to these criteria, if these patients had at least three features of the minor and three major, they were recognized as patients with atopic dermatitis.

Major criteria includes pruritus, flexor area dermatitis, chronic and recurrent dermatitis, general family history of atopy, asthma, allergic rhinitis and atopic dermatitis, and minor criteria includes dry skin, actinosis vulgaris, keratosis pilaris, pityriasis alba, Anterior neck folds, dani morgan folds, periorbital darkness, recurrent conjunctivitis, Increase of palmar lines, hand dermatitis, foot dermatitis, facial erythema, pale face, susceptibility to skin infections, white dermatographism, lips dermatitis, intolerance of wool and plastic, intolerance to solvents and detergents, increased itching with sweating, exacerbation of the disease with stimuli, foods and stress (25).

The control group was healthy women without atopic dermatitis. After selecting subjects and controls using Hanifin and Rajka criteria, the demographic and Premenstrual Symptoms Screening Test (PSST) questionnaires were completed by both groups. The demographic questionnaire included questions such as age, education, occupation, marital status, number of pregnancies, and questions about menstruation. The PSST questionnaire was consisted of 14 questions and had two parts:

The first part consists of 9 symptoms of mood and behavior, and the second part is the effect of these symptoms on people's lives and includes 5 questions. The Likert questionnaire is based on 4 options (at all, mild, moderate and severe), scoring from 0 to 3. In order to diagnose moderate or severe premenstrual syndrome (PMS), there must be three conditions: option 1 to 4 is at least one moderate or severe, in addition to the previous one, option 1 to 9 is at least 4 moderate or

severe, and at five last question, there is a moderate or severe case. The minimum score is zero and the maximum score is 42 points. Validity and reliability of the questionnaire were also confirmed (26).

Finally, the collected data were analyzed using SPSS-21 software and statistical tests of Kolmogorov-Smirnov, T-independent, Mann-Whitney and Chi-square were analyzed. $P < 0.05$ was considered significant.

Results

The mean age of participants in the dermatitis group was 30.17 ± 7.45 in the case group without dermatitis (27.29 ± 8.03) ($p = 0.04$). The majority of subjects in the case group were housewives (53.7%) and the majority of the control group were students (43.9%) ($p = 0.004$) (Table 1).

Table 1. Distribution of demographic and midwifery variables in two groups of atopic dermatitis and without atopic dermatitis (N=41)

Variables	Case mean \pm SD	Control mean \pm SD	P-value
Age (year)	30.17 \pm 7.45	27.29 \pm 8.03	0.04**
Age of first menstruation (year),	13.74 \pm 1.62	14 \pm 1.97	0.52*
Duration of bleeding (day),	6.59 \pm 3.03	5.63 \pm 1.81	0.13**
Periodic cycle (day),	26.64(4.45)	27.46(3.28)	0.34**
occupation	N(%)	N(%)	
Employed	14(34.1)	6(14.6)	0.004
housewife	22(53.7)	17(41.5)	
Student	5(12.3)	18(43.9)	
education .			
Under diploma	13(31.7)	11(26.8)	0.67
diploma	6(14.6)	9(22)	
Academic	22(53.7)	21(51.2)	
Marital status			
married	29(7.07)	21(51.2)	0.07
single	12(29.3)	20(48.8)	
Gravida			
0	16(39)	22(53.7)	0.35
1	11(26.8)	10(24.4)	
≥ 2	14(34.1)	9(9.22)	

* Independent T test, ** Mann-Whitney test, rest: Kai-two

The results of single-variable analysis of the relationship between atopic dermatitis and severity of premenstrual syndrome showed that the risk of moderate or severe premenstrual syndrome in subjects with atopic dermatitis was 7.9% more than healthy subjects (CI-95%: 2.57-21.58, OR=7.49). Because of the significant difference in some of the variables in the

studied groups and in order to control the possible interventional factors, variables with a significant level of less than 0.1 were re-entered into logistic regression. By controlling these factors, the probability of having moderate or severe premenstrual syndrome in subjects with atopic dermatitis was 13.26 times more than healthy subjects (CI-95%:3.31-53.07, OR=13.26) (Table 2).

Table 2. Comparison of severity of premenstrual syndrome in case and control group

Variables	Severity of premenstrual syndrome		Unadjusted		Adjusted*	
	Moderate to severe N=29	Without or mild N=53	OR(CI-95%)	p-value	OR(C-95%)	p-value
Atopic dermatitis N(%)						
+	23(56.1)	18(43.9)	7.49(2.75-21.58)	<0.0001	13.26(3.31-53.07)	<0.0001
-	6(14.6)	35(85.4)	1		1	

* adjusted for age, job, marriage

Discussion

The findings of this study showed that there is a significant relationship between the severity of premenstrual syndrome and atopic dermatitis, so that in patients with atopic dermatitis, the probability of having moderate or severe premenstrual syndrome is 13.26 times that of healthy people. The results of this study indicate that there is a relationship between hormonal effects and clinical manifestations of atopic dermatitis. The results of other studies are also consistent with the results of our study.

In some studies, it has been suggested that hormonal factors have been effective in dermatosis in people with dermatitis, and atopic dermatitis is more common in adult women (8, 7). Some studies have shown that patients with eczema are more susceptible to rheumatoid arthritis than normal controls (27, 10). In some studies, worsening skin symptoms have been shown during menstruation.

The prevalence of menstrual cycles, which is associated with worsening symptoms of atopic dermatitis, varies from 9% to 100% in these studies (28). Kemmett et al. In their study stated that there may be a weak link between the symptoms of atopic dermatitis in premenstruation and the presence of PMS symptoms (29).

In a study by Kiriya et al., PMS symptoms were worse in all patients with dermatitis during this period, while these symptoms were not found in people whose skin lesions were not worse during this period (19). In two studies, it has been shown that skin reactivity to

anti-allergenic substances and anti-gluing agents increases during premenstrual phase (31, 30), and psychosomatic conditions such as premenstrual syndrome accounts for more than half of the cases relate to anxiety and depression and eczema (1).

Considering that the variables of age, occupation and marriage in the study group in either of the two groups had a significant level of 0.1 or less, they could be considered as possible interventional factors. Prior to controlling these factors, moderate to severe PMS in individuals with atopic dermatitis was about 7.5 times higher, which increased to 13.26 times after controlling for these variables, which is significant. It seems that further studies are needed in this area so that more definitive results can be obtained.

One of the limitations of this study is the low prevalence of atopic dermatitis and the difficult and time-consuming access to patients, which required a long time to collect data, and good collaboration among participants in the study was one of the strengths of this study.

The results showed that the incidence of moderate to severe menstrual syndrome in atopic dermatitis was more than 13 times higher than healthy subjects. The results of this study further clarify the relationship between atopic dermatitis and premenstrual syndrome. Considering the strong relationship between these two variables, it is suggested that in the presence of premenstrual syndrome, the possibility of atopic dermatitis should be considered and treated if necessary.

Acknowledgment

Heraby, we would like to thank the Vice-Chancellor for Research and Technology of Hamedan University of

Medical Sciences for financing the project, as well as the officials and personnel of the medical clinic of Sina Hospital and all the participants in this study.

References

1. Yarcheski A, Mahon NE, Yarcheski TJ. Anger in early adolescent boys and girls with health manifestations. *Nurs Res*. 2002;51(4):229-36.
2. Gupta MA, Gupta AK. Psychiatric and psychological co-morbidity in patients with dermatologic disorders. *Am J Clin Dermatol*. 2003;4(12):833-42.
3. Masoumi Z, Shayan A, Ahobeiri f, Roshanaei G, Khanialamooti M. The effect of training premenstrual coping skills in undergraduate students in the school of nursing and midwifery, hamadan-iran. *Prevent Care Nurs Midwife J*. 2017;7(1):44-51. [In Persian].
4. Gupta MA, Gupta AK, Ellis CN, Koblenzer CS. Psychiatric evaluation of the dermatology patient. *Dermatolog Clin*. 2005;23(4):591-9.
5. Masoumi SZ, Alamoti MK, Shobeiri F, Roshanaei G, Mohaghahi H. Evaluating the prevalence of premenstrual syndrome among female undergraduate students of School of Nursing and Midwifery, Hamadan university of medical sciences in Iran. 2016. 831-7.
6. Radmanesh MsS. Psychosomatic skin diseases. *Iran J Dermatol*. 1997;1(3-4):7-34.
7. Chen W, Mempel M, Schober W, Behrendt H, Ring J. Gender difference, sex hormones, and immediate type hypersensitivity reactions. *Allergy*. 2008;63(11):1418-27.
8. Kornizky Y, Topilsky M, Fireman E, Kivity S, Kivity S. Specific IgE antibodies to aeroallergens and food among Israelis. *Ann Allergy Asthma Immunolog*. 1999;83(2):149-52.
9. Montnemery P, Nihlén U, Löfdahl CG, Nyberg P, Svensson Åk. Prevalence of self-reported eczema in relation to living environment, socio-economic status and respiratory symptoms assessed in a questionnaire study. *BMC dermatol*. 2003;3(1):4.
10. Koo J, Lebwohl A. Psychodermatology: The Mind and Skin Connection. *Am family physic*. 2001;64(11).
11. Richard BO, William DJ, Timothy GB. Andrews' diseases of the skin: clinical dermatology. *Clin dermatol*. 2000;10:641-43.
12. MacKie RM, Champion RH, Burton JL, Burns DA, Breathnach SM. Textbook of dermatology. Textbook of dermatology. 1998.
13. Jordaan HF. The diagnosis and management of eczema in pregnancy: review article. *Curr Aller Clin Immunol*. 2009;22(1):18-22.
14. Fölster-Holst R, Pape M, Buss YL, Christophers E, Weichenthal M. Low prevalence of the intrinsic form of atopic dermatitis among adult patients. *Allergy*. 2006;61(5):629-32.
15. Ingordo V, D'Andria G, D'Andria C, Tortora A. Results of atopy patch tests with house dust mites in adults with intrinsic and extrinsic atopic dermatitis. *J Eur Acad Dermatol Venereol*. 2002;16(5):450-4.
16. Wüthrich B, Schmid-Grendelmeier P. The atopic eczema/dermatitis syndrome. Epidemiology, natural course, and immunology of the IgE-associated (" extrinsic") and the nonallergic (" intrinsic") AEDS. *J Investigat Allergol Clin Immunol*. 2002;13(1):1-5.
17. Kemmett D, Tidman MJ. The influence of the menstrual cycle and pregnancy on atopic dermatitis. *Brit J Dermatol*. 1991;125(1):59-61.
18. Weatherhead S, Robson SC, Reynolds NJ. Eczema in pregnancy. *Brit Med J*. 2007;335(7611):152.
19. Kiriyaama K, Sugiura H, Uehara M. Premenstrual deterioration of skin symptoms in female patients with atopic dermatitis. *Dermatol*. 2003;206(2):110-2.
20. Timonen M, Jokelainen J, Hakko H, Silvennoinen-Kassinen S, Meyer-Rochow VB, Herva A, et al. Atopy and depression: results from the Northern Finland 1966 birth cohort study. *Mol Psychiatry*. 2003;8(8):738-44.

21. Timonen M, Jokelainen J, Silvennoinen-Kassinen S, Herva A, Zitting P, Xu B, et al. Association between skin test diagnosed atopy and professionally diagnosed depression: a Northern Finland 1966 Birth Cohort study. *Biological psychiatry*. 2002;52(4):349-55.
22. Kempainen T, Mäkityrö J, Jokelainen P, Nieminen M, Järvelin M. Is grand multiparity associated with offsprings' hospital-treated mental disorders? A 28-year follow-up of the North Finland 1966 birth cohort. *Soc Psychiat Psychiat Epidemiol*. 2000;35(3):104-8.
23. Pekkanen J, Xu B, Järvelin MR. Gestational age and occurrence of atopy at age 31—a prospective birth cohort study in Finland. *Clin Experiment Allerg*. 2001;31(1):95-102.
24. Kljakovic M, Pullon S. Allergy and the premenstrual syndrome (PMS). *Allergy*. 1997;52(6):681-3.
25. Joji T. Diagnostic standard for atopic dermatitis. *JMA Policies*. 2002;45(11):460-5.
26. Armand A, Talaee A. Investigating the Efficacy of Cognitive-Behavioral Stress-Management Training on Decreasing the Psychological Problems and Symptoms of Premenstrual Syndrome of Afflicted Women. *Iran J Obste, Gynecol Infertil*. 2012;15(21):24-7.
27. Hashiro M, Okumura M. Anxiety, depression and psychosomatic symptoms in patients with atopic dermatitis: comparison with normal controls and among groups of different degrees of severity. *J Dermatol Sci*. 1997;14(1):63-7.
28. Cho S, Kim HJ, Oh SH, Park CO, Jung JY, Lee KH. The Influence of Pregnancy and Menstruation on the Deterioration of Atopic Dermatitis Symptoms. *Ann Dermatol*. 2010;22(2):180-5.
29. Agner T, Damm P, Skouby SO. Menstrual cycle and skin reactivity. *J Am Acad Dermatol*. 1991;24(4):566-70.
30. Alexander S. Patch testing and menstruation. *Lancet (London, England)*. 1988;2(8613):751.